OptiCleanse™ GHI Sugar- & Stevia-Free‡

Support for Gastrointestinal System, Hepatic Function, and Cytokine Balance[†]



Available in Vanilla Delight

‡This formula is not a low-calorie dietary supplement. Please see the Supplement Facts panel for more details.

Discussion

OptiCleanse™ GHI Sugar- & Stevia-Free[‡] is now sweetened with a natural, high-potency sweetener extracted from monk fruit. This generally recognized as safe (GRAS) monk-fruit extract offers a high-quality sweetness and flavor without the bitter aftertaste associated with some natural sweeteners.

Like XYMOGEN's other OptiCleanse GHI formulas, OptiCleanse™ GHI Sugar- & Stevia-Free[‡] contains macro- and micronutrients, as well as a host of ingredients (some patented or proprietary) that support fatty acid metabolism, gastrointestinal health, and healthy eicosanoid and cytokine metabolism. Activated cofactors support mitochondrial energy production needed for biotransformation and detoxification. This formula's ingredients help moderate phase I detoxification, upregulate and support phase II pathways, and provide antioxidant support as well.[†]

Protein Metabolism

VegaPro is XYMOGEN's proprietary blend of pea protein isolate and rice protein concentrate, L-glutamine, glycine, and taurine. Generation of glutathione and sulfation cofactors—vital for phase II conjugation—requires an array of amino acids. The combination of pea protein and rice protein, containing a complement of amino acids, achieves an amino acid score of 100%. Glutamine, a conditionally essential and versatile amino acid with two nitrogen moieties, is crucial to nitrogen metabolism and helps maintain healthy liver tissue and function.^[1,2] The amino acid glycine is needed for bile synthesis, phase II detoxification, and glutathione production. Taurine, a derivative of the sulfur-containing amino acid cysteine, is also important for synthesis of bile salts and helps stabilize cell membranes.[†]

Gastrointestinal Support

Ginger root, included to support healthy digestion such as the release of bile from the gallbladder, acts at several sites to moderate PGE(2) production and support the normal response to inflammation. Fiber (from inulin and flaxseeds) supports production of short-chain fatty acids as well as a healthy intestinal flora.

MeadowPure™, an organic flaxseed complex, possesses excellent oxidative stability, supports antioxidant activity, and provides lignins, soluble fiber, and omega-3 and omega-6 essential fatty acids. The flaxseed is extensively milled, which provides a smoother mouthfeel and less grit than earlier versions of OptiCleanse formulas. Glutamine plays a key role in healthy intestinal cell proliferation and gut barrier integrity, immune function, and normal tissue healing.

11.1.2

Clinical Applications

- » Supports Natural Detoxification Mechanisms[†]
- » Supports Gastrointestinal Health[†]
- » Supports a Balanced Cytokine Profile[†]
- » Lactose-Free Vegan Protein Source†

OptiCleanse™ GHI Sugar- & Stevia-Free‡ is a comprehensive, monkfruit-extract-sweetened, low-allergy—potential dietary supplement designed to support gastrointestinal (GI) function and balanced detoxification. It features VegaPro™, XYMOGEN's proprietary amino acid and pea/rice protein blend; Aminogen®, to facilitate protein absorption; phytonutrients; mineral amino acid chelates; and activated B vitamins, including Quatrefolic® and methylcobalamin. In conjunction with a modified elimination diet, OptiCleanse GHI Sugar- and Stevia-Free‡ addresses GI and hepatic function as well as eicosanoid balance and cytokine metabolism. This formula is suitable for vegans.†

Detoxification Support

Ellagic acid (from pomegranate extract) prevents over-induction of CYP1A enzymes, works at the gene level to induce synthesis of glutathione-S-transferases and other phase II activities, binds directly to toxins, and protects DNA and hepatocytes. [5,6] Watercress is a rich source of beta-phenylethyl isothiocyanate (PEITC)—a versatile compound found to inhibit phase I enzymes and induce the phase II enzymes associated with biotransformation and excretion of toxins. Watercress has been found to contain even stronger phase II inducers known as 7-methylsulfinyheptyl and 8-methylsulfinyloctyl isothiocyanates as well. [7,8] Green tea catechins not only support antioxidant activity but also appear to act as modulators of phase I and phase II detoxification. [9] Choline is present to support lipid metabolism in the liver and can be converted to betaine, a methyl donor. fi10]

The active, bioavailable form of **B vitamins** (pyridoxal-5'-phosphate (B6), 5-methyltetrahydrofolate (folate), methylcobalamin (B12)) and glycine all support amino acid conjugation and are vital for the detoxification of xenobiotics and xenoestrogens. 5-methyltetrahydrofolate (5-MTHF), methylcobalamin, betaine, and **methylsulfonylmethane** (MSM) are present to support methylation and detoxification. 5-MTHF supports healthy folate nutrition, especially in individuals with variations in folate metabolism. In OptiCleanse GHI, 5-MTHF is provided as Quatrefolic® for enhanced stability, solubility, and bioavailability.^{f[11]}

Preventium®, a patented form of potassium hydrogen d-glucarate, supports glucuronidation. Sulfation is supported by **MSM** and **sodium sulfate**. Acetylation is supported by **d-calcium pantothenate**, pyridoxal-5'-phosphate, and magnesium. Several minerals in OptiCleanse GHI are provided as Albion® mineral chelates and TRAACS® mineral amino acid chelates for enhanced gastrointestinal absorption and bioavailability.^{112]}

Antioxidant Support and Cytokine Balance

Bioflavonoids, quercetin, rutin, and curcumin support antioxidant activity, counter free radicals, and support healthy eicosanoid and cytokine metabolism. [13,14] Curcumin has a long history of use for its support of a normal, healthy response to inflammation. [15] N-acetyl-cysteine (NAC) stimulates glutathione synthesis, enhances glutathione-S-transferase activity, and promotes detoxification. [16] Selenium glycinate provides support for glutathione metabolism and antioxidant protection. †

OptiCleanse™ GHI Sugar- & Stevia-Free Vanilla Delight Supplement Facts

Serving Size: 2 Scoops (about 54 g)

	%Daily Value
	14%*
	15%*
	4%3
	20%*
	**
	52%
2500 IU	50%
250 mg	417%
15 mg	1000%
5 mg	294%
40 mg	200%
5 mg	250%
200 mcg	50%
50 mcg	833%
150 mcg	50%
35 mg	350%
240 mg	24%
6 mg	33%
60 mcg	40%
140 mg	35%
10 mg	67%
100 mcg	143%
2 mg	100%
60 mcg	50%
35 mcg	47%
500 mg	21%
510 mg	15%
5.6 g	*
1.28 g	*:
392 mg	*
400 mg	*:
250 mg	*
250 mg	*
250 mg	
250 mg	*
200 mg	*:
200 mg	*:
150 mg	*
150 mg	*:
120 mg	*:
100 mg	*
100 mg	*:
100 mg	*:
82 mg	**

Other Ingredients: VegaPro™ (XYMOGEN's proprietary blend of pea protein concentrate, pea protein isolate, taurine, glycine, rice protein concentrate, and L-glutamine), sunflower oil, natural flavors (no MSG), medium-chain triglyceride oil, tripotassium citrate, cellulose gum, xanthan gum, Aminogen®, monk fruit extract, guar gum, and silica.

DIRECTIONS: Blend, shake, or briskly stir 2 level scoops (54 g) into 10-12 ounces of chilled, pure water (or mix amount for desired thickness) and consume once daily, or use as directed by your healthcare practitioner.

Consult your healthcare practitioner prior to use. Individuals taking medication should discuss potential interactions with their healthcare practitioner. Do not use if tamper seal is damaged. The labeling on this product does not comply with California's Proposition 65. Therefore, this product may not be sold in California.

STORAGE: Keep closed in a cool, dry place out of reach of children.

DOES NOT CONTAIN: Wheat, gluten, yeast, soy, animal or dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, or artificial preservatives.







References

- Smith RJ, Wilmore DW. Glutamine nutrition and requirements. JPEN J Parenter Enteral Nutr. 1990 Jul-Aug;14(4 Suppl):94S-99S. Review. [PMID: 2119461]
- Lacey JM, Wilmore DW. Is glutamine a conditionally essential amino acid? Nutr Rev. 1990 Aug;48(8):297-309. Review. [PMID: 2080048]
- Lantz RC, Chen GJ, Sarihan M, et al. The effect of extracts from ginger rhizome on inflammatory mediator production. *Phytomedicine*. 2007 Feb;14(2-3):123-28. [PMID: 16709450]
- Adolphe JL, Whiting SJ, Juurlink BH, Thorpe LU, Alcorn J. Health effects with consumption of the flax lignan secoisolariciresinol diglucoside. *Br J Nutr.* 2010 Apr;103(7):929-38. Review. [PMID: 20003621]
- Barch DH, Rundhaugen LM, Stoner GD, et al. Structure-function relationships of the dietary anticarcinogen ellagic acid. *Carcinogenesis*. 1996 Feb;17(2):265-9. [PMID: 8625448]
- Girish C, Koner BC, Jayanthi S, et al. Hepatoprotective activity of picroliv, curcumin and ellagic acid compared to silymarin on paracetamol induced liver toxicity in mice. Fundam Clin Pharmacol. 2009 Dec;23(6):735-45. [PMID: 19656205]
- Rose P, Faulkner K, Williamson G, et al. 7-Methylsulfinylheptyl and 8-methylsulfinyloctyl isothiocyanates from watercress are potent inducers of phase II enzymes. *Carcinogenesis*. 2000 Nov;21(11):1983-8. [PMID: 11062158]
- B. Hofmann T, Kuhnert A, Schubert A, et al. Modulation of detoxification enzymes by watercress: in vitro and in vivo investigations in human peripheral blood cells. Eur J Nutr. 2009 Dec;48(8):483-91. [PMID: 19636603]
- Akhlaghi M, Bandy B. Dietary green tea extract increases phase 2 enzyme activities in protecting against myocardial ischemia-reperfusion. *Nutr Res.* 2010 Jan;30(1):32-39. [PMID: 20116658]
- Linus Pauling Institute. http://lpi.oregonstate.edu/infocenter/othernuts/choline/. Accessed May 8, 2012.
- 11. Quatrefolic. http://www.quatrefolic.com/. Accessed May 8, 2012.
- 12. Albion. http://www.albionminerals.com/. Accessed May 8, 2012.
- 13. Garg R, Gupta S, Maru GB. Dietary curcumin modulates transcriptional regulators of phase I and phase II enzymes in benzo[a]pyrene-treated mice: mechanism of its anti-initiating action. *Carcinogenesis*. 2008 May;29(5):1022-32. [PMID: 18321868]
- Amália PM, Possa MN, Augusto MC, et al. Quercetin prevents oxidative stress in cirrhotic rats. *Dig Dis Sci.* 2007 Oct;52(10):2616-21. [PMID: 17431769]
- Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of Curcuma longa: a review of preclinical and clinical research. *Altern Med Rev.* 2009 Jun;14(2):141-53. [PMID: 19594223]
- Kelly GS. Clinical applications of N-acetylcysteine. Altern Med Rev. 1998 Apr;3(2):114-27. Review. [PMID: 9577247]

Additional references available upon request

Typical Amino Acid Profile Per Serving:

Alanine	1,129 mg
Arginine	2,260 mg
Aspartic Acid	3,049 mg
Cysteine	248 mg
Glutamic Acid	4,450 mg
Glycine	1,578 mg
Histidine	652 mg
Isoleucine	1,214 mg
Leucine	2,208 mg
Lysine	1,910 mg
Methionine	279 mg
Phenylalanine	1,432 mg
Proline	1,141 mg
Serine	1,404 mg
Taurine	981 mg
Threonine	500 mg
Tryptophan	258 mg
Tyrosine	1,005 mg
Valine	1,317 mg

Aminogen® is a registered trademark of Triarco Industries. Aminogen® is protected under US patent 5,387,422. Preventium® is a registered trademark of Applied Food Sciences, LLC. (US patents 4,845,123, 5,364,644, 5,561,160).